

ON CONJUGACY CLASSES AND DERIVED LENGTH

EDITH ADAN-BANTE

ABSTRACT. Let G be a finite group and A , B and D be conjugacy classes of G with $D \subseteq AB = \{xy \mid x \in A, y \in B\}$. Denote by $\eta(AB)$ the number of distinct conjugacy classes such that AB is the union of those. Set $\mathbf{C}_G(A) = \{g \in G \mid x^g = x \text{ for all } x \in A\}$. If $AB = D$ then $\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))$ is an abelian group. If, in addition, G is supersolvable, then the derived length of $\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))$ is bounded above by $2\eta(AB)$.

1. INTRODUCTION

Let G be a finite group, $A \subset G$ be a conjugacy class of G , i.e. $A = a^G = \{g^{-1}ag \mid g \in G\}$ for some a in G . Let X be a normal subset of G , i.e. $X^g = \{g^{-1}xg \mid x \in X\} = X$ for all $g \in G$. We can check that X is a union of n distinct conjugacy classes of G , for some integer $n > 0$. Set $\eta(X) = n$.

We can check that given any two conjugacy classes A and B of G , the product $AB = \{xy \mid x \in A, y \in B\}$ of A and B is a normal subset of G . Then $\eta(AB)$ is the number of distinct conjugacy classes of G such that AB is the union of those classes.

Denote by $\mathbf{C}_G(X) = \{g \in G \mid x^g = x \text{ for all } x \in X\}$ the centralizer of X in G . If G is a solvable group, denote by $\text{dl}(G)$ the derived length of G . Let $\mathbf{Z}(G)$ be the center of the group G .

In this note, we are exploring the relations between the structure of the group G and the product AB of some conjugacy classes A and B of G . More specifically, we are exploring the relation between the derived length of some section of G and properties of AB .

Given a finite solvable group G and conjugacy classes A and B of G , is there any relationship between the derived length of G and $\eta(AB)$? In general, the answer seems to be no. For instance $A\{e\} = A$ for any finite group G and any conjugacy class A of G . Thus $\eta(AB)$ may not give us information about $\text{dl}(G)$, but it does give us a linear bound on the derived length of a section of G , namely on the section $\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))$.

Theorem A. *Let G be a finite group and A , B be normal subsets of G . Then $\mathbf{C}_G(AB)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))$ is abelian.*

Given any integer $m > 0$, we show in Example 2.1 that there exists a nilpotent group G with conjugacy classes A , B and D such that $|\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))| = m$ and $AB = D$. It follows that although $\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))$ is abelian, its order is unbounded and even the number of distinct prime divisors is unbounded.

Date: 2009.

1991 Mathematics Subject Classification. 20d15.

Key words and phrases. Conjugacy classes, solvable groups, derived length, products of characters, irreducible constituents.

Theorem B. *Let G be a supersolvable group, A , B and D conjugacy classes of G such that $D \subseteq AB$. Then $\text{dl}(\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))) \leq 2\eta(AB)$.*

We now mention the “dual” situation for characters, where the “dual” of a conjugacy class A is an irreducible character χ and the “dual” of the Kernel $\text{Ker}(\chi)$ of χ is the centralizer $\mathbf{C}_G(A)$ of A . Let Ξ and Ψ be complex characters of G . Since any product of characters is a character, $\Xi\Psi$ is a character of G . Thus it can be written as an integral linear combination of irreducible characters of G . Let $\eta(\Xi\Psi)$ be the number of distinct irreducible constituents of the character $\Xi\Psi$. In Theorem A of [4], it is proved that there exist universal constants c and d such that for any solvable group G , any irreducible characters χ, ψ and θ such that θ is a constituent of $\chi\psi$, we have that $\text{dl}(\text{Ker}(\theta)/(\text{Ker}(\chi) \cap \text{Ker}(\psi))) \leq c\eta(\chi\psi) + d$.

In Theorem A of [7] is proved that given any supersolvable group G and any conjugacy class A , we have that $\text{dl}(G/\mathbf{C}_G(A)) \leq 2\eta(AA^{-1}) - 1$. We conjecture in [7] that there exist universal constants q and r such that for any solvable group G and any conjugacy class A of G , we had that $\text{dl}(G/\mathbf{C}_G(A)) \leq q\eta(AA^{-1}) + r$. In light of Theorem B and because of the “dual” situation with characters, namely Theorem A of [4], we wonder the following

Conjecture C. *There exist universal constants r and s such that for any solvable group G , any conjugacy classes A , B and D of G such that $D \subseteq AB$, we have that*

$$\text{dl}(\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))) \leq r\eta(AB) + s.$$

We will show in Theorem 2.3 that the previous conjecture has an affirmative answer if and only if Conjecture of [7] has an affirmative answer. We would like to point out that there are several examples of “dual results” between products of conjugacy classes and products of character. For example, see [9], [2] and [8], [3] and [5]. However not every result in products of characters has a “dual” result in conjugacy classes, see for instance Section 3.

In Section 4 we provide an example of a property in a conjugacy class A of G that bounds the nilpotent class of $G/\mathbf{C}_G(A)$, and therefore it bounds the derived length of that section.

2. PROOFS

Notation. Let G be a group, X be a subset of G and $a \in G$. Set $[a, X] = \{[a, x] \mid x \in X\}$. Observe that $a^X = \{a[a, x] \mid x \in X\} = a[a, X]$.

Proof of Theorem A. Write $N = \mathbf{C}_G(AB)$ and $C = \mathbf{C}_G(N)$ so $AB \subseteq C$. Let $a \in A$, $b \in B$ and $g \in C$. Then ab and ab^g lie in AB , and so lie in C , and since C is a subgroup, it follows that $b^{-1}b^g$ also lies in C . It follows then than working in G/C , b is central, so $[\langle b \rangle, G] \subseteq C$ and $[\langle b \rangle, G, N] = 1$. In particular, $[\langle b \rangle, N, N] = 1$ and so by the three-subgroups lemma, $[N', \langle b \rangle] = 1$. Since this holds for all $b \in B$, we have that $N' \subseteq \mathbf{C}_G(B)$. Similarly, $N' \subseteq \mathbf{C}_G(A)$ and the result follows. $\square$

Example 2.1. Let $m > 0$ be an integer. Write $m = \prod_{i=1}^l p_i$, where p_i are primes not necessarily distinct. Let P_i be nonabelian of order p_i^3 , and let G be the direct product of the groups P_i . Choose noncommuting elements a_i and b_i in P_i and write $a = \prod a_i$ and $b = \prod b_i$. It is easy to check that $A = a^G = aZ$, $B = b^G = bZ$, where $Z = \mathbf{Z}(G)$. Then $AB = abZ = D$, where $D = (ab)^G$. Also $\mathbf{C}_G(A) \cap \mathbf{C}_G(B) = Z$, which has order $\prod p_i$, while $\mathbf{C}_G(D)$ has order $\prod (p_i)^2$. Then $|\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))| = \prod_{i=1}^l p_i = m$, as wanted.

Definition 2.2. Let $F(n)$ be a nondecreasing function defined on the natural numbers. A group G is **good** for F if for every conjugacy class A of G , the group $G/\mathbf{C}_G(A)$ is solvable with derived length at most $F(\eta(AA^{-1}))$.

In Theorem A of [7], it is proved that $F(n) = 2n - 1$ is good for all supersolvable groups.

Theorem 2.3. *Suppose that a function F is good for all homomorphic images of G . Let A, B and D be conjugacy classes of G with $D \subseteq AB$. Then $\mathbf{C}_G(D)/(\mathbf{C}_G(A) \cap \mathbf{C}_G(B))$ is solvable with derived length at most $1 + F(\eta(AB))$.*

Proof. Let $N = \mathbf{C}_G(D)$, and observe that $\mathbf{C}_G(A) \cap \mathbf{C}_G(B) \subseteq N$. It suffices to show that both $N/(N \cap \mathbf{C}_G(A))$ and $N/(N \cap \mathbf{C}_G(B))$ have derived length at most $1 + F(\eta(AB))$. We will prove the required length inequality for $N/(N \cap \mathbf{C}_G(A))$; the other inequality follows similarly.

Let $C = \mathbf{C}_G(N)$ and, using the standard bar convention, write $\bar{G} = G/C$. Since $D \subseteq C$, we see that the identity is an element of $\bar{A}\bar{B}$. Also $\bar{A}$ and $\bar{B}$ are conjugacy classes of $\bar{G}$, so they must be inverse classes. By Lemma 2.5 of [5] we have that $\eta(\bar{A}\bar{B}) \leq \eta(AB)$, and so by hypothesis we have then

$$(2.4) \quad \text{dl}(\bar{G}/\mathbf{C}_{\bar{G}}(\bar{A})) \leq F(\eta(\bar{A}\bar{B})) \leq F(\eta(AB)).$$

Thus $\text{dl}(G/K) = F(\eta(\bar{A}\bar{A}^{-1}))$, where K is the preimage in G of $\mathbf{C}_{\bar{G}}(\bar{A})$. It follows then that $\text{dl}(N/(N \cap K)) \leq F(\eta(\bar{A}\bar{A}^{-1}))$. Now $\mathbf{C}_G(A) \subseteq K$ and so it will be enough to show that $N \cap K/(N \cap \mathbf{C}_G(A))$ is abelian, yielding the desired derived length bound for $N/(N \cap \mathbf{C}_G(A))$.

Now K centralizes A modulo C , so $[\langle a \rangle, K] \subseteq C$ and $[\langle a \rangle, K, N] = 1$ for $a \in A$. Then $[\langle a \rangle, (N \cap K), (N \cap K)] = 1$ and by the three-subgroups lemma, $(N \cap K)'$ centralizes a , and we have $(N \cap K)' \subseteq N \cap \mathbf{C}_G(A)$ as wanted. $\square$

Since $F(n) = 2n - 1$ is good for supersolvable groups (Theorem A of [7]), Theorem A follows from the previous result.

3. CONJUGACY CLASS SIZES

Fix a prime p . Let G be a finite p -group and A be conjugacy classes of G . In Theorem A of [5], we proved that if $|A| = p^n$ for some integer n , then $\eta(AA^{-1}) \geq n(p - 1) + 1$. Thus, in the particular case that G is p -group, there is a relation between the size of A and $\eta(AA^{-1})$. We want to point out a “dual” result in character theory, where the “dual” of a conjugacy class is an irreducible complex character χ and the “dual” of the inverse of a conjugacy class is the complex conjugate character $\bar{\chi}$ of χ , where $\bar{\chi}(g) = \overline{\chi(g)}$ for all $g \in G$. More specifically, in Theorem A of [3] is proved that if G is a p -group, χ is an irreducible character of G and $\chi(1) = p^n$, then the product $\chi\bar{\chi}$ of χ and $\bar{\chi}$ has at least $2n(p - 1) + 1$ distinct irreducible constituents, i.e $\eta(\chi\bar{\chi}) \geq 2n(p - 1) + 1$.

In Theorem A of [1] is proved that if χ is an irreducible character of a solvable group G with $\chi(1) > 1$, then $\chi(1)$ has at most $\eta(\chi\bar{\chi}) - 1$ different prime factors. If, in addition, G is supersolvable, then $\chi(1)$ has at most $\eta(\chi\bar{\chi}) - 2$ prime factors. Is there any “dual” result in conjugacy classes as Theorem A of [1] in characters? In other words,

Question 3.1. *Does it exist a function $f : \mathbb{Z} \rightarrow \mathbb{Z}$ such that for any solvable group G and any conjugacy class A of G , we have that $|A|$ has at most $f(\eta(AA^{-1}))$ different prime factors?*

The answer is no, such function can not exist. More specifically,

Example 3.2. Let p be a prime and P be a group of order p . Let G be the group of order $p(p-1)$, where $P \triangleleft G$ and G/P induces on P the full group of automorphisms of P . Then P contains just one nontrivial conjugacy class A of G , namely $A = P \setminus \{1\}$. Observe that $AA^{-1} = P$ and so $\eta(AA^{-1}) = 2$. Also $P = \mathbf{C}_G(A)$, and thus $|G/\mathbf{C}_G(A)| = p-1$. This is obviously unboundedly large, and by a result of Erdos [10], it has unboundedly many prime factors.

Remark. Let G be the group as in the previous Example. Let λ be an irreducible character of N . Then λ is a linear character and the induced character λ^G is an irreducible character of degree $p-1$. Set $\chi = \lambda^G$. We can check then $\eta(\chi\bar{\chi}) = p$, namely the irreducible constituents of $\chi\bar{\chi}$ are χ and all irreducible character with kernel containing N . Since $n-1 \leq 2^{n-2}$ for any integer $n > 0$, then $p-1$ has at most $p-2$ distinct prime factors.

4. CONJUGACY CLASSES AND NILPOTENT CLASS

Let $\mathbf{Z}_1(G) = \mathbf{Z}(G)$ be the center of the group G and by induction define the i -center of G as $\mathbf{Z}_i(G)/\mathbf{Z}_{i-1}(G) = \mathbf{Z}(G/\mathbf{Z}_{i-1}(G))$. The following is a well known result.

Lemma 4.1. *Let N be a group. Write $Z_m = Z_m(N)$ for the m -th center of N and write N^m for the m -th term of the lower central series of N . Then $[N^m, Z_m] = 1$.*

Proof. Induct on m . For $m = 1$, we have $[N^1, Z_1] = [N, \mathbf{Z}(N)] = 1$, as needed. For $m > 1$, we want $[N^{m-1}, N, Z_m] = 1$. We have $[N, Z_m, N^{m-1}] \subseteq [Z_{m-1}, N^{m-1}] = 1$ by the inductive hypothesis. Now work in $\bar{N} = N/Z$ where $Z = Z_1 = \mathbf{Z}(N)$. Note that $\overline{N^{m-1}} = (\bar{N})^{m-1}$ and $\overline{Z_m} = Z_{m-1}(\bar{N})$. Then

$$1 = [Z_{m-1}(\bar{N}), (\bar{N})^{m-1}] = [\overline{Z_m}, \overline{N^{m-1}}] = [\overline{Z_m}, \overline{N^{m-1}}],$$

and we have $[Z_m, N^{m-1}] \subseteq Z$. Then $[Z_m, N^{m-1}, N] = 1$, and the result follows by the three-subgroups lemma. $\square$

Theorem 4.2. *Let $N \trianglelefteq G$ and $a \in G$. Assume that $[N, a] \subseteq Z_m$, where $Z_m = Z_m(G)$ is the m -th center of N . Let $C = \mathbf{C}_N(a^N)$. Then N/C is nilpotent of class at most m .*

Proof. Induct on m . if $m = 0$ we are assuming that $[N, a] = 1$ and so N centralizes all of a^N and $C = N$. In this case N/C is nilpotent of class zero. We can assume therefore that $m > 0$. Our goal is to show that $N^{m+1} \subseteq C$.

Now let $x \in a^N$. We want $[N^{m+1}, \langle x \rangle] = 1$, or equivalently, $[N^m, N, \langle x \rangle] = 1$. Now $[N, \langle x \rangle, N^m] \subseteq [Z_m, N^m] = 1$ by the previous lemma.

Let $Z = Z_1 = \mathbf{Z}(N)$ and write $\bar{G} = G/Z$. Then

$$Z_{m-1}(\bar{N}) = \overline{Z_m} \supseteq [\overline{N}, \bar{a}] = [\bar{N}, \bar{a}],$$

and thus by the inductive hypothesis, N/B is nilpotent of class at most $m-1$, where B is the preimage in N of $\mathbf{C}_{\bar{N}}((\bar{a})^{\bar{N}}) = \mathbf{C}_{\bar{N}}(\overline{a^N})$. In particular, $N^m \subseteq B$ so x centralizes N^m modulo Z , and we have $[\langle x \rangle, N^m] \subseteq Z$, and hence $[\langle x \rangle, N^m, N] = 1$. The three-subgroups lemma now yields $[N^m, N, \langle x \rangle] = 1$, as wanted. $\square$

Remark. Let $G_1 = C_2 \wr C_2$ be the wreath product of C_2 by C_2 , where C_2 is the cyclic group of order 2. Thus $|G_1| = 8$ and G_1 is non abelian. Let $a_1 \in G_1 \setminus \mathbf{Z}(G_1)$. We can check that $a_1^{G_1} = a\mathbf{Z}(G_1)$ and $G_1/\mathbf{C}_{G_1}(a_1^{G_1})$ is abelian and so it is nilpotent of class 1.

Let $N = G_1 \times G_1$ and $a_2 = (a_1, a_1)$ in N . Observe that C_2 acts on N by permuting the entries. Set $G_2 = C_2 N$. We can check that $a_2^{G_2} \subset a_2 \mathbf{Z}_2(G_2)$ but $a_2^{G_2} \not\subset a_2 \mathbf{Z}(G_2)$, and $G_2/(\mathbf{C}_{G_2}(a_2^{G_2}))$ is nilpotent of class 2.

The author wonders if given any integer $m > 2$, we can find a group G with an element $a \in G$ such that $a^G \subseteq a\mathbf{Z}_m(G)$ and $G/\mathbf{C}_G(a^G)$ is nilpotent of class m .

Acknowledgment. I thank Harvey Blau and Michael Bush for useful discussions and emails. I would like to thank the Instituto de Matematicas, UNAM, unidad Morelia, and Gerardo Raggi for their hospitality while visiting. Also, I thank the referee of a previous version of this note for helpful comments and suggestions of how to improve both the presentation and the results of this note; those are included here with her/his permission.

REFERENCES

- [1] E. Adan-Bante, Products of characters and derived length, J. of Algebra, 266 (2003), 305-319.
- [2] E. Adan-Bante, Products of characters and finite p -group, J. of Algebra, 277 (2004), 236-255.
- [3] E. Adan-Bante, Products of characters and finite p -groups II, Archiv der Mathematik, 82 No 4 (2004), 289-297.
- [4] E. Adan-Bante, Products of characters and derived length II, J. Group Theory, 8 (2005), 453-459.
- [5] E. Adan-Bante, Conjugacy classes and finite p -groups, Arch. Math. 85 (2005) 297-303.
- [6] E. Adan-Bante, Homogeneous products of conjugacy classes, Arch. Math. 86 (2006) 289-294.
- [7] E. Adan-Bante, Derived length and products of conjugacy classes, Israel J. Math. 168 (2008), 93-100.
- [8] E. Adan-Bante, On nilpotent groups and conjugacy classes, to appear Houston Journal of Mathematics.
- [9] Z. Arad, E. Fisman, An analogy between products of two conjugacy classes and products of two irreducible characters in finite groups, Proc. of the Edinburgh Math. Soc. 30 (1987), 7-22.
- [10] P. Erdos, On the normal number of prime factors of $p-1$ and some related problems concerning Euler's ϕ -function, Q. J. Math., Oxf. Ser. 6, 205-213 (1935).

DEPARTMENT OF MATHEMATICAL SCIENCE, NORTHERN ILLINOIS UNIVERSITY, WATSON HALL
320 DEKALB, IL 60115-2888, USA

E-mail address: EdithAdan@illinoisalumni.org